



Synthelis®

GPCR

Protein Catalog

CCR2a - C-C Chemokine Receptor type 2 isoform A

PL022

Product specification

Acronym: CCR2a

Synonyms: MCP1R

Origin species : Human

Protein reference : P41597 (UniProtKB)
U03882.1 (GenBank)

Family: GPCR class A

Expression system: E.coli based CFPS

Format: Proteoliposomes

Protein sequence: Met1 – Ala374

Tag : 6xHis tag (N-ter)

Cleavage site: Factor Xa

Product MW: 44.5 kDa

Application: Drug screening & discovery, antibody development, structural biology

Product description

CCR2 is a member of the beta chemokines receptors family. This gene encodes two isoforms of a receptor for monocyte chemoattractant protein-1, a chemokine which specifically mediates monocyte chemotaxis. Monocyte chemoattractant protein-1 is involved in monocyte infiltration in inflammatory diseases such as rheumatoid arthritis as well as in the inflammatory response against tumor. The receptors encoded by this gene mediate agonist-dependent calcium release.

Recombinant protein sequence

His tag – factor X cleavage site -

MLSTSRSRFIRNTNESGEEVTTFFDYDYGAPCHKFDVKQIGAQLLPPLYSLVFIFGFVGNMLVVLILINCKKLKCLTDIYLLNLAISDLL
FLITLPLWAHSAANEWVFGNAMCKLFTGLYHIGYFGGIFFIILLTIDRYLAIVHAVFALKARTVTFGVVTSVITWLVAVEASVPGIIFT
KCQKEDSVYVCGPYFPRGWNNFHTIMRNILGLVPLLMVICYSGILKTLRLCRNEKKRHRAVRVIFTIMIVYFLFWTPYNIVILLNT
FQEFFGLSNCESTSQLDQATQVTETLGMTHCCINPIIYAFVGEKFRSLFHIALGCRIAPLQKPVCGGPGVVRPGKNVKVTTQGLLDGR
GKGKSIGRAPEASLQDKEGA



Quality analysis

Purity:

Liposomes are directly incorporated into the Cell-Free reaction, thus, some impurities from the *E.coli* lysate might be present in the proteoliposomes.

A negative control (proteoliposomes without the protein of interest) can be provided (useful for screening, immunization...).

The purity can be improved by protein expression in detergent and relipidification after purification step(s).

Purification procedure: CCR2a proteoliposomes are purified on a sucrose gradient.

NB : Migration of membrane proteins on SDS-PAGE can results in « gel shifting » due to the presence of hairpins (helix-loop-helix)¹⁻³.

References :

1 – Rath A., et al., Detergent binding explains anomalous SD-PAGE migration of membrane proteins PNAS, 2009 Feb 10, vol. 106

2 – Rath A., et al., Acrylamide concentration determines the direction and magnitude of helical membrane protein gel shifts, PNAS, 2013 Sep 24, 110(39)

3 – Rath A., et al., Correction factors for membrane protein molecular weight readouts on sodium dodecyl sulfate-polyacrilamide gel electrophoresis, Anal. Biochem., 2013 Mar 1, 434(1)

Formulation

Buffer: Available in Hepes 50mM, pH 7.5 with cryoprotectants. Other buffers or customized formulation can be provided upon request.

Customized Hydrophobic matrix: Customized formulation with specific lipids like PEGylated or biotinylated lipids can be used upon request, as well as targeting molecules.

Storage/Stability: Store at +4°C for up to one week or several months at -80°C. Aliquot for storage.
Do not freeze-thaw after aliquoting.

Use restrictions: For life science research use only.

Available sizes: 10 µg, 50 µg, 100 µg, customized quantity on request.





Fig.1: Identification of CCR2a in the proteoliposomes by Western Blot (using an anti-6xHis antibody).



Need a specific amount, a quote or any additional information?
Contact-us



Synthelisis

T : +33 (0)4 76 54 95 35
E: contact@synthelisis.fr
www.synthelisis.com

